arXiv:1601.03291v1 [math.GR] 12 Jan 2016

ON THE SUBGROUPS AND NORMAL SUBGROUPS OF THE GROUP REPRESENTATION OF THE CAYLEY TREE.

F. H. HAYDAROV

Abstract. In this paper, we give a characterization of the normal subgroups of index $2^s(2n+1)$, $s \in \{1,2\}$, $n \in \mathbb{N}$ and of the subgroups of index three of the group representation of the Cayley tree.

Key words. G_k- group, subgroup, normal subgroup, homomorphism, epimorphism.

AMS Subject Classification: 20B07, 20E06.

1. Introduction

In the theory of groups there are very important unsolved problems, most of which arise in solving of problems of natural sciences as physics, biology etc. In particular, if configuration of a physical system is located on a lattice (in our case on the graph of a group) then the configuration can be considered as a function defined on the lattice. Usually, more important configuration (functions) are the periodic ones. It is well-known that if the lattice has a group representation then periodicity of a function can be defined by a given subgroup of the representation. More precisely, if a subgroup, say H, is given, then one can define a H- periodic function as a function, which has a constant value (depending only on the coset) on each (right or left) coset of H. So the periodicity is related to a special partition of the group (that presents the lattice on which our physical system is located). There are many works devoted to several kinds of partitions of groups (lattices) (see e.g. [1], [3], [5], [7]).

One of the central problems in the theory of Gibbs measures is to study periodic Gibbs measures corresponding to a given Hamiltonian of model. For any subgroup H of the group G_k we define H-periodic Gibbs measures. To find new periodic and weakly periodic Gibbs measures one usually needs to find new subgroups of the group representation of the Cayley tree. In Chapter 1 of [5] it is given a one to one correspondence between the set of vertices V of the Cayley tree Γ^k and the group G_k and it is given a full description of subgroups of index two and there are also constructed several normal subgroups of the group G_k. But there wasn't a full description of normal subgroups of finite index (without index two) and to the best of our knowledge until now there wasn't any description of a not normal subgroup of finite index of the group representation of the Cayley tree. In [4] it is given a full description of normal subgroups of indices 4 and 6 for the group representation of the Cayley tree. In this paper we continue this investigation and construct all normal subgroups of index $2^s(2n+1)$, $s \in \{1,2\}$, $n \in \mathbb{N}$ and all subgroups of index three for the

group representation of the Cayley tree.

Cayley tree and its group representation. A Cayley tree (Bethe lattice) Γ^k of order $k \geq 1$ is an infinite homogeneous tree, i.e., a graph without cycles, such that exactly $k+1$ edges originate from each vertex. Let $\Gamma^k = (V, L)$ where V is the set of vertices and L that of edges (arcs). Let G_k be a free product of $k+1$ cyclic groups of the second order with generators $a_1, a_2, ...a_{k+1}$, respectively. It is known that there exists a one to one correspondence between the set of vertices V of the Cayley tree Γ^k and the group G_k. To give this correspondence we fix an arbitrary element $x_0 \in V$ and let it correspond to the unit element e of the group G_k. Using $a_1, ..., a_{k+1}$ we label the nearest-neighbors of element e, moving in positive direction. Now we'll label the nearest-neighbors of each $a_i, i = 1, ..., k+1$ by $a_i a_j, j = 1, ..., k+1$. Since all a_i have the common neighbor e we have $a_i a_i = a_i^2 = e$. Other neighbors are labeled starting from $a_i a_i$ in positive direction. We label the set of all the nearest-neighbors of each $a_i a_j$ by words $a_i a_j a_q, q = 1, ..., k+1$ starting from $a_i a_j a_j = a_i$ by the positive direction. Iterating this argument one gets a one-to-one correspondence between the set of vertices V of the Cayley tree Γ^k and the group G_k.

Any(minimal represented) element $x \in G_k$ has the following form: $x = a_{i_1} a_{i_2} ... a_{i_n}$, where $1 \leq i_m \leq k+1, m = 1, ..., n$. The number n is called the length of the word x and is denoted by $l(x)$. The number of letters $a_i, i = 1, ..., k+1$, that enter the non-contractible representation of the word x is denoted by $w_x(a_i)$.

The following result is well-known in group theory. If φ is a homomorphism of a group G with the kernel H then H is a normal subgroup of the group G and $\varphi(G) \simeq G/H$, (where G/H is the quotient group) so the index $|G : H|$ coincides with the order $|\varphi(G)|$ of the group $\varphi(G)$.

Usually we define natural homomorphism g from G onto the quotient group G/H by the formula $g(a) = aH$ for all $a \in G$. Then $Ker\varphi = H$.

Definition 1. *Let $M_1, M_2, ..., M_m$ be some sets and $M_i \neq M_j$, for $i \neq j$. We call the intersection $\cap_{i=1}^m M_i$ contractible if there exists $i_0 (1 \leq i_0 \leq m)$ such that*

$$\cap_{i=1}^m M_i = \left(\cap_{i=1}^{i_0-1} M_i\right) \cap \left(\cap_{i=i_0+1}^m M_i\right).$$

Let $N_k = \{1, ..., k+1\}$. The following Proposition describes several normal subgroups of G_k.

Put

$$H_A = \left\{ x \in G_k \ \mid \ \sum_{i \in A} \omega_x(a_i) \ is \ even \right\}, \quad A \subset N_k. \tag{1.1}$$

Proposition 1. *[5] For any $\emptyset \neq A \subseteq N_k$, the set $H_A \subset G_k$ satisfies the following properties:*
(a) H_A is a normal subgroup and $|G_k : H_A| = 2$;
(b) $H_A \neq H_B$, for all $A \neq B \subseteq N_k$;
(c) Let $A_1, A_2, ..., A_m \subseteq N_k$. If $\cap_{i=1}^m H_{A_i}$ is non-contractible, then it is a normal subgroup of index 2^m.

Theorem 1. *[5]*

1. *The group G_k does not have normal subgroups of odd index ($\neq 1$).*
2. *The group G_k has normal subgroups of arbitrary even index.*

2. Subgroups and normal subgroups of finite index.

2.1. Normal subgroups of index $\mathbf{2^s(2n+1)}$, $\mathbf{s \in \{1,2\}}$, $\mathbf{n \in \mathbb{N}}$.

Definition 2. *A commutative group G of order 2^n is called $K_{2^n}-$group if G is generated by free product of n elements $c_i, i = \overline{1,n}$ satisfying the relations $o(c_i) = 2,\ i = \overline{1,n}$,*

Proposition 2. *Let φ be a homomorphism of the group G_k onto a finite commutative group G. Then $\varphi(G_k)$ is isomorphic to K_{2^i} for some $i \in \mathbb{N}$.*

Proof. Let $(G, *)$ be a commutative group of order n and $\varphi : G_k \to G$ be an epimorphism. We will first show that $n \in \{2^i |\ i \in \mathbb{N}\}$. Suppose n does not belong to $\{2^i |\ i \in \mathbb{N}\}$. Then there exist $m > 1,\ m, s \in \mathbb{N} \cup \{0\}$ such that $n = 2^s m$, where m is odd. Since $\varphi : G_k \to G$ is an epimorphism there exist distinct elements $\varphi(a_{j_p}),\ p = \overline{1,s}$. Clearly, $< \varphi(a_{j_1}), \varphi(a_{j_2})..., \varphi(a_{j_s}) >$ is a subgroup of G and $| < \varphi(a_{j_1}), \varphi(a_{j_2})..., \varphi(a_{j_s}) > | = 2^s$. Since $m > 1$ there exist at least one element $a_{j_0} \in G_k$ such that $\varphi(a_{j_0}) \neq \varphi(a_{j_i}),\ i \in \{1, 2, ..., s\}$. Hence $< \varphi(a_{j_0}), \varphi(a_{j_1})..., \varphi(a_{j_s}) >$ is a subgroup of G. Then by Lagrange's theorem $|G| : 2^{s+1}$ which contradicts our assumption that m is odd. Thus $n \in \{2^i |\ i \in \mathbb{N}\}$. If $n = 2^q$ then G is equal to $< \varphi(a_{j_1}), \varphi(a_{j_2})..., \varphi(a_{j_q}) >$ We define the following mapping $\gamma(a_{j_i}) = c_i \in K_q$ for all $i \in \{1, 2, ..., q\}$. It is easy to check that this mapping is a one to one correspondence from G to K_q. This completes the proof. □

Let $A_1, A_2, ..., A_m \subset N_k,\ m \in \mathbb{N}$ and $\cap_{i=1}^m H_{A_i}$ is non-contractible. Then we denote by $\Re$ the following set

$$\Re = \{\cap_{i=1}^m H_{A_i} |\ A_1, A_2, ..., A_m \subset N_k,\ m \in \mathbb{N}\}.$$

Theorem 2. *Let φ be a homomorphism from G_k to a finite commutative group. Then there exists an element H of $\Re$ such that $Ker\varphi \simeq H$ and conversely.*

Proof. We give a one to one correspondence between $\{Ker\varphi |\ \varphi(G_k) \simeq G \in \{K_{2^i} |\ i \in \mathbb{N}\}\}$ and $\Re$. Let φ be a homomorphism from G_k to a finite commutative group of order p. Then by Proposition 2 the number p belongs to the set $\{2^i |\ i \in \mathbb{N}\}$ and if $p = 2^n$ then $\varphi : G_k \to K_{2^n}$ is an isomorphism. For any nonempty sets $A_1, A_2, ..., A_n \subset N_k$ and $a_i \in G_k,\ i \in N_k$ we define the following homomorphism

$$\phi_{A_1A_2...A_n}(a_i)=\begin{cases} c_j, \ \text{ if } \ i\in A_j\setminus(A_2\cup A_3\cup ...\cup A_n),\ j=\overline{1,n}\\ .\ .\ .\ .\ .\ .\ .\ .\\ c_{j_1}c_{j_2}, \ \text{ if } \ i\in(A_{j_1}\cap A_{j_2})\setminus(A_1\cup ...A_{j_1-1}\cup A_{j_1+1}\cup ...\\ ...\cup A_{j_2-1}\cup A_{j_2+1}\cup ...\cup A_n),\ 1\le j_1<j_2\le n\\ .\ .\ .\ .\ .\ .\ .\ .\\ c_1...c_{j-1}c_{j+1}...c_n, \ \text{ if } \ i\in(A_1\cap ...\cap A_{j-1}\cap A_{j+1}...\cap A_n)\setminus A_j,\ j=\overline{1,n}\\ c_1c_2...c_n, \ \text{ if } \ i\in(A_1\cap A_2\cap ...\cap A_n)\\ e, \ \text{ if } \ i\in N_k\setminus(A_1\cup A_2\cup ...\cup A_n). \end{cases}$$

If $i\in\emptyset$ then we'll accept that there is no index $i\in N_k$ satisfying the required condition. For $x\in G_k$ if $\varphi_{A_1A_2...A_n}(x)=e$ the number of $a_i,\ i=\overline{1,n}$ of a_i appearing in the word x must be even. Therefore

$$Ker\varphi_{A_1A_2...A_n}=H_{A_1}\cap H_{A_2}\cap ...\cap H_{A_n}.$$

Thus the following equality holds (up to isomorphism)

$$\{Ker\varphi|\ \varphi(G_k)\simeq G\in\{K_{2^i}|\ i\in\mathbb{N}\}\}=\Re.$$

□

The group G has finitely generators of the order two. Assume that r is the minimal number of such generators of the group G and without loss of generality we can take these generators to be $b_1,b_2,...b_r$. Let e_1 be the identity element of the group G. We define a homomorphism from G_k onto G. Let $\Xi_n=\{A_1,\ A_2,...,A_n\}$ be a partition of $N_k\setminus A_0,\ 0\le|A_0|\le k+1-n$. Then we consider the homomorphism $u_n:\{a_1,a_2,...,a_{k+1}\}\to\{e_1,b_1...,b_m\}$ given by

$$u_n(x)=\begin{cases} e_1, \ \text{ if } \ x=a_i, i\in A_0\\ b_j, \ \text{ if } \ x=a_i, i\in A_j, j=\overline{1,n}. \end{cases}\tag{2.1}$$

For $b\in G$ we denote by $R_b[b_1,b_2,...,b_m]$ a representation of the word b by generators $b_1,b_2,...,b_r,\ r\le m$. Define the homomorphism $\gamma_n:G\to G$ by the formula

$$\gamma_n(x)=\begin{cases} e_1, \ \text{ if } \ x=e_1\\ b_i, \ \text{ if } \ x=b_i, i=\overline{1,r}\\ R_{b_i}[b_1,...,b_r], \ \text{ if } \ x=b_i,\ i\notin\{1,...,r\}. \end{cases}\tag{2.2}$$

Put

$$H^{(p)}_{\Xi_n}(G)=\{x\in G_k|\ l(\gamma_n(u_n(x))):2p\},\ 2\le n\le k-1.\tag{2.3}$$

Let $\gamma_n(u_n(x)))=\tilde{x}$. We introduce the following equivalence relation on the set G_k : $x\sim y$ if $\tilde{x}=\tilde{y}$. It's easy to check this relation is reflexive, symmetric and transitive.

Let $(G,*)$ be a group of order n, generated by two elements of order 2, and let $\Im_n$ be the set of all such groups.

Proposition 3. *For the group G_k the following equality holds*

$$\{Ker\varphi|\ \varphi: G_k \to G \in \Im_{2n} \text{ is an epimorphism}\} =$$
$$= \{H^{(n)}_{B_0B_1B_2}(G)|\ B_1, B_2 \text{ is a partition of the set } N_k \setminus B_0,\ 0 \le |B_0| \le k-1\}.$$

Proof. For an arbitrary group $G \in \Im_{2n}$ we give a one to one correspondence between the two given sets. Let e_1 be the identity element of G and B_1, B_2 be a partition of $N_k \setminus B_0$, where $B_0 \subset N_k,\ 0 \le |B_0| \le k-1$. Then we define the homomorphism $\varphi_{B_0B_1B_2}: G_k \to G$ by the formula

$$\varphi_{B_0B_1B_2}(a_i) = \begin{cases} b_1, & \text{if } \ i \in B_1 \\ b_2, & \text{if } \ i \in B_2. \end{cases} \tag{2.4}$$

There is only one such a homomorphism (corresponding to B_0, B_1, B_2). It's clearly $x \in Ker\varphi_{B_0B_1B_2}$ if and only if $\tilde{x}$ is equal to e_1. Hence it is sufficient to show that if $y \in H^{(n)}_{B_0B_1B_2}(G)$ then $\tilde{y} = e_1$. Suppose that there exist $y \in G_k$ such that $l(\tilde{y}) \ge 2n$. Let $\tilde{y} = b_{i_1}b_{i_2}...b_{i_q},\ q \ge 2n$ and $S = \{b_{i_1}, b_{i_1}b_{i_2}, ..., b_{i_1}b_{i_2}...b_{i_q}\}$. Since $S \subseteq G$ there exist $x_1, x_2 \in S$ such that $x_1 = x_2$ which contradicts the fact that $\tilde{y}$ is a non-contractible. Thus we have proved that $l(\tilde{y}) < 2n$. Since $y \in H^{(n)}_{B_0B_1B_2}(G)$ the number $l(\tilde{y})$ must be divisible by $2n$. Thus for any $y \in H^{(n)}_{B_0B_1B_2}(G)$ we have $\tilde{y} = e_1$. Hence for the group G we get $Ker\varphi_{B_0B_1B_2} = H^{(n)}_{B_0B_1B_2}(G)$. □

Let us denote by $\aleph_n$ the following set

$$\{H^{(n)}_{B_0B_1B_2}(G)|\ B_1, B_2 \text{ is a partition of the set } N_k \setminus B_0,\ 0 \le |B_0| \le k-1,\ |G| = 2n\}.$$

Theorem 3. *Any normal subgroup of index $2n$ with $n = 2^i(2s+1),\ i = \{0,1\},\ s \in \mathbb{N}$ has the form $H^{(n)}_{B_0B_1B_2}(G),\ |G| = 2n$ i.e.,*

$$\aleph_n = \{H|\ H \lhd G_k,\ |G_k : H| = 2n\}$$

Proof. We will first prove that $\aleph_n \subseteq \{H|\ H \lhd G_k,\ |G_k : H| = 2n\}$. Let G be a group with $|G| = 2n$ and B_1, B_2 a partition of the set $N_k \setminus B_0,\ 0 \le |B_0| \le k-1$. For $x = a_{i_1}a_{i_2}...a_{i_n} \in G_k$ it's sufficient to show that $x^{-1}H^{(n)}_{B_0B_1B_2}(G)\ x \subseteq H^{(n)}_{B_0B_1B_2}(G)$. We have (as in the proof of Proposition 3) if $y \in H^{(n)}_{B_0B_1B_2}(G)$ then $\tilde{y} = e_1$, where e_1 is the identity element of G. Now we take any element z from the set $x^{-1}H^{(n)}_{B_0B_1B_2}(G)\ x$. Then $z = x^{-1}h\ x$ for some $h \in H^{(n)}_{B_0B_1B_2}(G)$. We have

$$\tilde{z} = \gamma_n(v_n(z)) = \gamma_n\left(v_n(x^{-1}h\ x)\right) = \gamma_n\left(v_n(x^{-1})v_n(h)v_n(x)\right) =$$
$$= \gamma_n\left(v_n(x^{-1})\right)\gamma_n\left(v_n(h)\right)\gamma_n\left(v_n(x)\right) = \left(\gamma_n\left(v_n(x)\right)\right)^{-1}\gamma_n\left(v_n(h)\right)\gamma_n\left(v_n(x)\right).$$

From $\gamma_n(v_n(h)) = e_1$ we get $\tilde{z} = e_1$ i.e., $z \in H^{(n)}_{B_0B_1B_2}(G)$. Hence

$$H^{(n)}_{B_0B_1B_2}(G) \in \{H|\ H \lhd G_k,\ |G_k : H| = 2n\}.$$

Now we'll prove that $\{H|\ H \lhd G_k,\ |G_k : H| = 2n\} \subseteq \aleph_n$. Let $H \lhd G_k,\ |G_k : H| = 2n$. We consider the *natural homomorphism* $\phi : G_k \to G_k : H$ i.e. the homomorphism given by $\phi(x) = xH,\ x \in G_k$. There exist $e, b_1, b_2, ...b_{2n-1}$ such that $\phi : G_k \to \{H, b_1H, ..., b_{2n-1}H\}$ is an epimorphism. Let $(\{H, b_1H, ..., b_{2n-1}H\}, *) = \wp$, i.e., $\wp$ is the factor group. If we show that $\wp \in \Im_{2n}$ then the theorem will be proved. Assuming that $\wp \notin \Im_{2n}$ then there are $c_1, c_2, ..., c_q \in \wp,\ q \geq 3$ such that $\wp =< c_1, ..., c_q >$. Clearly $< c_1, c_2 >$ is a subgroup of $\wp$ and $| < c_1, c_2 > |$ is greater than three.

`Case` **n=2s+1.** By Lagrange's theorem $| < c_1, c_2 > | \in \left\{m|\ \frac{2n}{m} \in \mathbb{N}\right\}$. If e_2 is the identity element of $\wp$ then from $c_1^2 = e_2$ we take $| < c_1, c_2 > | = 2n$. Hence $< c_1, c_2 >= \wp$ but $c_3 \notin < c_1, c_2 >$. Thus $\wp \in \Im_{2n}$.

`Case` **n=2(2s+1).** From Lagrange's theorem $| < c_1, c_2 > | \in \{4, 2(2s+1), 4(2s+1)\}$. Let $| < c_1, c_2 > | = 4$. If the number four isn't equal to one of these numbers $| < c_1, c_3 > |$ or $| < c_2, c_3 > |$ then we'll choose that pairs. If $| < c_1, c_2 > | = | < c_1, c_3 > | = | < c_2, c_3 > | = 4$ then $< c_1, c_2, c_3 >$ is a group of order eight. Again by Lagrange's theorem $|\wp| = 2n$ must be divisible by 8 which is impossible.

Let $| < c_1, c_2 > | = 2(2s+1)$. Then

$$< c_1, c_2 >= \{e,\ c_1,\ c_2,\ c_1c_2,\ c_1c_2c_1,\ \ldots,\ \underbrace{c_1c_2...c_1}_{4s}\} = A.$$

It's easy to check that

$$c_3A \cup A \subseteq \wp, \qquad c_3A \cap A = \emptyset, \qquad |c_3A \cup A| = |c_3A| + |A| = 2n = |\wp|.$$

We then deduce that $c_3A \cup A = \wp$ but we'll show $c_3c_1c_3 \in \wp$ does not belong to $c_3A \cup A$. Since c_1, c_2, c_3 are generators we have $c_3c_1c_3 \notin < c_1, c_2 >$ so $c_3c_1c_3 = xc_3$ with $x \in < c_1, c_2 >$. But $x = c_3c_1 \notin < c_1, c_2 >$.

If $| < c_1, c_2 > | = 4(2l+1)$ then $< c_1, c_2 >= \wp$ but $c_3 \notin < c_1, c_2 >$. Hence $\wp \in \Im_{2n}$. This completes the proof. □

2.2. **Subgroups of index three.** In this section we will characterize the subgroups of index three for the group representation of the Cayley tree.

Let $(A_1,\ A_2)$ be a partition of the set $N_k \backslash A_0,\ 0 \leq |A_0| \leq k-1$. Put m_j be a minimal element of $A_j,\ \ j \in \{1, 2\}$. Then we consider the function $u_{A_1A_2} : \{a_1, a_2, ..., a_{k+1}\} \to \{e, a_1..., a_{k+1}\}$ given by

$$u_{A_1A_2}(x) = \begin{cases} e, & \text{if } \ x = a_i,\ i \in N_k \setminus (A_1 \cup A_2) \\ a_{m_j}, & \text{if } \ x = a_i,\ i \in A_j,\ j = \overline{1,2}. \end{cases}$$

Define $\gamma :< e, a_{m_1}, a_{m_2} > \to \{e, a_{m_1}, a_{m_2}\}$ by the formula

$$\gamma(x) = \begin{cases} e \text{ if } \ x = e \\ a_{m_1} \text{ if } \ x \in \{a_{m_1},\ a_{m_2}a_{m_1}\} \\ a_{m_2} \text{ if } \ x \in \{a_{m_2},\ a_{m_1}a_{m_2}\} \\ \gamma\left(a_{m_i}a_{m_{3-i}}a_{m_i}...\gamma(a_{m_i}a_{m_{3-i}})\right) \text{ if } \ x = a_{m_i}a_{m_{3-i}}a_{m_i}...a_{m_{3-i}},\ l(x) \geq 3,\ i = \overline{1,2} \\ \gamma\left(a_{m_i}a_{m_{3-i}}a_{m_i}...\gamma(a_{m_{3-i}}a_{m_i})\right) \text{ if } \ x = a_{m_i}a_{m_{3-i}}a_{m_i}...a_{m_i},\ l(x) \geq 3,\ i = \overline{1,2} \end{cases}$$

For a partition $(A_1,\ A_2)$ of the set $N_k \backslash A_0,\ 0 \leq |A_0| \leq k-1$ we consider the following set.

$$\Sigma_{A_1A_2}(G_k) = \{x \in G_k |\ \gamma(u_{A_1A_2}(x)) = e\}$$

Lemma 1. *Let $(A_1,\ A_2)$ be a partition of the set $N_k \backslash A_0,\ 0 \leq |A_0| \leq k-1$. Then $x \in \sum_{A_1A_2}(G_k)$ if and only if the number $l(u_{A_1A_2}(x))$ is divisible by 3.*

Proof. Let $x = a_{i_1}a_{i_2}...a_{i_n} \in G_k$ and $l(u_{A_1A_2}(x))$ be odd (the even case is similar). Then we can write $u_{A_1A_2}(x) = a_{m_i}a_{m_{3-i}}...a_{m_i}a_{m_{3-i}}a_{m_i}$. $\gamma(u_{A_1A_2}(x))$ is equal to

$$\gamma(a_{m_i}a_{m_{3-i}}...a_{m_i}a_{m_{3-i}}a_{m_i}) = \gamma\left(a_{m_i}a_{m_{3-i}}...a_{m_i}\gamma(a_{m_{3-i}}a_{m_i})\right) = \gamma(\underbrace{a_{m_i}a_{m_{3-i}}...a_{m_{3-i}}a_{m_i}}_{l(u_{A_1A_2}(x))-3}) = ...$$

We continue this process until the length of the word $\gamma(u_{A_1A_2}(x))$ will be less than three. From $\gamma(u_{A_1A_2}(x)) = e$ we deduce that $l(u_{A_1A_2}(x))$ is divisible by 3.

Conversely if $l(u_{A_1A_2}(x))$ is divisible by 3 then x is generated by the elements $a_{m_1}a_{m_2}a_{m_1}$ and $a_{m_2}a_{m_1}a_{m_2}$. Since $\gamma(a_{m_1}a_{m_2}a_{m_1}) = \gamma(a_{m_2}a_{m_1}a_{m_2}) = e$ we get $x \in \sum_{A_1A_2}(G_k)$. □

Proposition 4. *For the group G_k the following equality holds*

$$\{K |\ K \text{ is a subgroup of } G_k \text{ of index } 3\} =$$

$$= \{\Sigma_{A_1A_2}(G_k) |\ A_1,\ A_2 \text{ is a partition of } N_k \setminus A_0\}.$$

Proof. Let K be a subgroup of the group G_k with $|G_k : K| = 3$. Then there exist $p, q \in N_k$ such that $|\{K,\ a_pK,\ a_qK\}| = 3$. Put

$$A_0 = \{i \in N_k |\ a_i \in K\}, \quad A_1 = \{i \in N_k |\ a_ia_p \in K\}, \quad A_2 = \{i \in N_k |\ a_ia_q \in K\}$$

From $|G_k : K| = 3$ we conclude that $\{A_1, A_2\}$ is a partition of $N_k \setminus A_0$. Let m_i be a minimal element of $A_i,\ i = \overline{1,2}$. If we show $\sum_{A_1A_2}(G_k)$ is a subgroup of G_k (corresponding to K) then it'll be given one to one correspondence between given sets. For $x = a_{i_1}a_{i_2}...a_{i_n} \in \sum_{A_1A_2}(G_k),\ y = a_{j_1}a_{j_2}...a_{j_m} \in \sum_{A_1A_2}(G_k)$ it is sufficient to show that $xy^{-1} \in \sum_{A_1A_2}(G_k)$. Let $u_{A_1A_2}(x) = a_{m_i}a_{m_{3-i}}....a_{m_s},\ u_{A_1A_2}(y) = a_{m_j}a_{m_{3-j}}....a_{m_t}$, where

$(i,j,s,t) \in \{1,2\}^4$. Since $u_{A_1A_2}$ is a homomorphism we have

$$l\left(u_{A_1A_2}(xy^{-1})\right) = \begin{cases} |l\left(u_{A_1A_2}(x)\right) - l(u_{A_1A_2}(y^{-1}))\,|, & \text{if } \ s=t \\ l\left(u_{A_1A_2}(x)\right) + l(u_{A_1A_2}(y^{-1}))\,, & \text{if } \ s\neq t. \end{cases}$$

By Lemma 1 we see that $l\left(u_{A_1A_2}(x)\right)$ is divisible by 3 and $l\left(u_{A_1A_2}(y^{-1})\right) = l\left(\left(u_{A_1A_2}(y)\right)^{-1}\right)$ is also divisible by 3, so $l\left(u_{A_1A_2}(xy^{-1})\right)$ is a multiple of 3, which shows that $xy^{-1} \in \sum_{A_1A_2}(G_k)$. This completes the proof. □

References

[1] Cohen,D.E, Lyndon,R.C., (1963), *Free bases for normal subgroups of free groups*, Trans.Amer.Math.Soc., 108, pp. 526-537.

[2] D.S, Malik, J.N, Mordeson, M.K, Sen., (1997), *Fundamentals of Abstract Algebra*, McGraw-Hill Com.

[3] Ganikhodjaev, N.N, Rozikov, U.A., (1997), *Description of periodic extreme Gibbs measures of some lattice model on the Cayley tree*, Theor.Math.Phys. 111, pp. 480-486.

[4] U.A, Rozikov, F.H, Haydarov., (2014), *Normal subgroups of finite index for the group represantation of the Cayley tree*, TWMS Jour.Pure.Appl.Math. 5, pp. 234-240.

[5] U.A, Rozikov., (2013) *Gibbs measures on a Cayley tree*, World Sci. Pub, Singapore.

[6] Kurosh, A.G., (1953) *Group theory*, Akademic Verlag, Berlin.

[7] Young, J.W., (1927), *On the partitions of a group and the resulting classification*, Bull.Amer.Math.Soc., 33, pp.453-461.

F.H.Haydarov, National University of Uzbekistan, Tashkent, Uzbekistan.
E-mail address: haydarov_ imc@mail.ru.